

DDO spectroscopic survey of MOST variable stars^{*}

Theodor Pribulla^{1,2}, Slavek Rucinski¹, Rainer Kuschnig³,
Waldemar Ogłóza⁴, Bogumil Pilecki⁵

¹*David Dunlap Observatory, University of Toronto, P.O. Box 360, Richmond Hill, Ontario, L4C 4Y6, Canada*

²*Astronomical Institute, Slovak Academy of Sciences, 059 60 Tatranská Lomnica, Slovakia*

³*Institut für Astronomie, Universität Wien, Türkenschanzstrasse 17, A-1180 Wien, Austria*

⁴*Mt. Suhora Observatory of the Pedagogical University ul. Podchorążych 2, 30-084 Cracow, Poland*

⁵*Warsaw University Astronomical Observatory, Al. Ujazdowskie 4, 00-478 Warszawa, Poland*

Accepted 0000 Month 00. Received 0000 Month 00; in original form 2007 March 17

ABSTRACT

A spectroscopic support survey of 103 objects observed by the MOST satellite is presented; 96 are variable stars with 83 of them being new MOST variable-star detections or stars with variability types verified and/or modified on the basis of the MOST data. Analysis of 241 medium-resolution spectra using the broadening-functions formalism yielded radial velocities, projected rotational velocities (for 31 targets for which it was possible) and spectral type estimates. Seven new spectroscopic binaries were discovered; orbital solutions are given for two of them (HD 73709, and GSC 0814-0323). The visual binary HD 46180 was found to be composed of two close binary stars (eclipsing and non-eclipsing one) very probably forming a physical quadruple system.

Key words: stars: close binaries - stars: eclipsing binaries – stars: variable stars

1 INTRODUCTION

MOST (Microvariability & Oscillations of STars) is a microsatellite dedicated to detection and characterisation of variability of stars. It uses a 15-cm telescope which feeds a CCD photometer through a single, custom, broadband optical filter (350 – 700 nm). The pre-launch characteristics of the mission are described by Walker et al. (2003) and the initial post-launch performance by Matthews et al. (2004). Although the main goal of the mission was to study pulsations of stars, continuous photometry of selected fields led to discovery of many new variables of different types. Several new variables were found among guide stars used to accurately point the telescope. After the failure of the CCD used for the satellite stabilization in January 2006, the number of guide stars utilized for this purpose in the science CCD was increased to a few dozen per field which led to detection of many variable stars. Observations in a single, wide-band filter and small photometric amplitudes often prevented correct determination of variability type for newly discovered objects. The brightness range where spectroscopic information is poor or absent is typically within $8 < V < 11$.

While not all of some 150 initially selected targets have been observed in this program, we have decided to publish our results because the David Dunlap Observatory was

closed on 2008 July 2 and no longer functions. Because we have no access to a similar 2-metre class telescope and because the current data form a homogenous and uniform sample, this survey is published in the present form, as one of the last legacies of this venerable observatory.

The present paper gives for the survey targets (i) an estimate of the spectral type, (ii) radial velocity (for most of stars just one observation is available), (iii) the projected rotational velocity, for cases of $v \sin i > 30 \text{ km s}^{-1}$. The spectral type was estimated on the basis of the best template producing the integral of the broadening-function (BF) closest to unity (see Section 3). For a few stars previously known or discovered by us to be spectroscopic binaries, we present radial velocities for individual components and preliminary orbits. MOST photometry of the newly discovered eclipsing binaries will be presented in a separate paper. Analysis of variables (including eclipsing binaries) in the field of M67 is given in Pribulla et al. (2008). We note that many new variables are δ Scuti or γ Doradus pulsators, where estimates of the spectral type and the projected rotational velocity, $v \sin i$, are important data for the asteroseismology modelling.

2 SPECTROSCOPIC OBSERVATIONS

Spectroscopic observations were obtained using the slit spectrograph in the Cassegrain focus of 1.88m telescope of the David Dunlap Observatory (DDO). The spectra were taken

^{*} Based on spectroscopic data from the David Dunlap Observatory, University of Toronto

Table 1. Objects observed within the DDO survey and their characteristics.

HD/BD	GSC	Cluster	V	Sp. publ.	Sp BF	#	RV [km/s]	$v \sin i$ [km/s]	Multiplicity	Note
HD 965	4664-0318		9.09	A5p	A5:	3	0.7			roAp suspected, no apparent puls.
HD 6242	0022-0517		7.98	K2	K7(G)	1	36.7			LPV-known
HD 8801	2817-2326		6.42	A7m	F3:	3	3.4	58		
HD 9014	5276-0313		6.48	A9V	F4	4			SB2	dScut-new
HD 9289	5274-0240		9.71	A3	A9	1	10.1			roAp-known, spots-new
BD+37 410	2816-2068	NGC752	9.94	F4III	F5	2			SB1	eclBin-new
HD 11720	2816-1892	NGC752	8.09	K0III	K0(G)	2	20.0			RGpuls-new
BD+36 364	2816-2276	NGC752	10.44	F2III	F4	8			SB2	gDor-new
BD+37 431	2816-2273	NGC752	9.85	F2III	F4	4			SB2	gDor-new
HD 11885	2816-0111	NGC752	7.12	K1III	K2(G)	2	10.7			RGpuls-new
	2816-2061	NGC752	10.44	F2V	F4	1	4.5	130		gDor-new
	2816-2039	NGC752	10.90	F0V	F3	1	3.6	41		gDor-new
BD+36 373	2816-0432	NGC752	10.60	F2V	F2	1	5.7	45		eclBin-new
BD+37 440	2816-1973	NGC752	9.13	F1V	F2	3	1.9	39		gDor-new
BD+37 444	2816-2176	NGC752	9.62	F3V	F3	8			SB2	gDor-new
BD+37 445	2816-2225	NGC752	10.70	F2III	F7	1	-55.1	45		var-new
HD 12899	0636-0631		8.14	F0	F4	4	-6.7	141		dScut-new
	0636-0130		10.15		K0(G)	1	2.8			LPV-new ?
BD+13 336	0636-0976		9.30	A5	A9:	2	38.2	32		in vis. binary, dScut-new
HD 20790	0060-1499		8.90	F0	F4	1	62.4	97		gDor-new
	0060-0969		10.20	F8	K2:	1	15.7			gDor-new
HD 20884	0060-0509		7.47	K0	K1(G)	1	26.0			RGpuls-new
HD 21839	5296-0253		9.76	F5	F5	1	2.4	79		var ?
HD 21949	5296-0283		9.27	F0	F4(G?)	1	6.8			LPV-new
HD 21994	5296-0672		8.62	K0	K0(G)	1	36.2			RGpuls-new
HD 293785	4758-0092		9.29	K7	K2(G)	1	-44.8			LPV-new
HD 32704	4758-0066		8.66	G8V	G6-7	7			SB2	spots-new?
HD 258857	0158-1480	NGC2244	9.02	K2IV-V	K2	1	31.4			LPV-new
HD 258916	0158-0444	NGC2244	9.64	A5	F2	4			SB2	W UMa-known
HD 46105A	0158-2674	NGC2244	7.13	A1p	A1:	9			SB2	in vis binary, spots-new
HD 46105B	0158-2674	NGC2244	8.62		A7:	1	28.8			
HD 46180A	0154-2360	NGC2244	9.10	A3V	A3:	6			SB2	in vis binary, eclBin-new
HD 46180B	0154-2360	NGC2244	9.50		F4-5	4			SB2	in vis binary, eclBin-new
HD 46324	0154-1047	NGC2244	7.98	K5III	K2(G)	2	48.3			LPV-new
HD 46375	0154-0891	NGC2244	7.84	K1IV	K1	2	0.8			
HD 259696	0154-0435	NGC2244	8.85	K7		1	16.7			LPV-new
HD 46517	0154-0867	NGC2244	8.07	A7V	F0	2	41.7	175		dScut-gDor-new
HD 261230	0750-0863	NGC2264	9.39	A7V	F2	1	21.0	104		dScut/PMS-new?
HD 261355	0750-1713	NGC2264	8.23	F5-A0	F3	1	-10.4	44		PMS-new
HD 261387	0750-1597	NGC2264	10.60	A2V	A1	1	-8.8	127		PMS-new
HD 47554	0750-1471	NGC2264	7.86	F2V	F5	2	34.6	31		gDor-new
HD 261683	0746-1733	NGC2264	8.13	K5III	K3-4(G)	1	8.5			RGpuls-new?
HD 261783	0750-1163	NGC2264	8.28	K3II-III	K3(G)	1	17.3			LPV-new
HD 262014	0750-1525	NGC2264	9.95	A7V	F1-2	1	38.6	52		var-new?
HD 262500	0750-1227	NGC2264	9.96	A6V	F2	1	21.8	159		dScut-new
HD 48331	0750-0423	NGC2264	9.07	A0	F3	3	16.7			spots-new
BD+10 1246	0750-0455	NGC2264	8.37	M1		1	6.6			var-new
HD 263236	0750-0685	NGC2264	8.10		F4	1	41.2	38		gDor-new
HD 263551	0755-0575		9.68	A7	F3	2	23.4	67		dScut-new
HD 58714	0764-0548		6.73	K0	K2-3(G)	1	56.5			LPV-new
HD 61199	0187-1892		8.18	A3	F2	2			SB2-3	dScut-new
BD+05 1735	0187-1557		8.75	M2III		1	1.7			LPV-new
HD 61696	0187-1562		8.06	K0	K0(G)	1	25.2			RGpuls-new
BD+06 1752	0191-1302		10.16	F2	F4-5	2	-8.9	37		dScut-new
BD+05 1747	0187-1737		8.98	K2	K2(G)	1	26.9			RGpuls-new?
HD 73709	1395-2006	M44	7.31	F2III	F5:	14			SB1	in vis. binary, eclBin-new
BD+12 1913	0813-1617	M67	9.42	K2	K3(G)	1	33.1			RGpuls-new
BD+12 1918	0813-1152	M67	8.89	K0	K0(G)	1	39.5			RGpuls-new
BD+12 1919	0813-1032	M67	8.67	K5		1	32.5			RGpuls-LPV-new
BD+28 1653	1949-1584		8.99	K0	K2(G)	1	33.4			
	0814-1493	M67	9.68	K2III	K2(G)	1	33.1			RGpuls??
	0814-2331	M67	8.03	K4III	K2(G)	1	34.2			RGpuls
BD+12 1924	0814-2405	M67	9.25	K2		1	35.8			RGpuls-new
	0814-0889	M67	11.17	F4	F4	2			SB3	eclBin-known, Algol
HD 75638	0814-0601	M67	7.76	F0	F5	8			SB3	in vis. binary, eclBin-new
BD+12 1926	0814-1515	M67	9.47	K3III	K2(G)	1	34.1			RGpuls-new?
HD 75700	0814-0795	M67	6.73	K0	K1-2(G)	2	-2.0			RGpuls-new
HD 75717	0817-1563	M67	8.63	F2	F2	2	7.9	67		dScut-new
BD+12 1929	0814-0323	M67	8.88		F8-9	20			SB2	eclBin-new
BD+12 2159	0833-1094		8.98	K2	K1(G)	2	13.8			RGpuls-new?
HD 102083	0275-0271		8.59	A5e	F3-4	1	12.1	214		dScut-known
HD 103548	0273-0462		7.15	F2	F4-5(G)	2	3.9	32		LPV-new
HD 114839	1996-1075		8.46	Am	F4-5	2	-8.4	70		dScut-new
HD 116462	5547-0592		7.26	F0	F4	3	-1.8	48		gDor-new
BD+17 2663	1460-0878		9.64	M0		3	-10.4			LPV-new
BD+17 2667	1460-0976		9.95	F8	F5	4	-17.9			
BD+17 2670	1460-0532		9.82	K2	K2(G)	2	-48.7			

Table 1. (continued)

HD/BD	GSC	Cluster	V	Sp. publ.	Sp BF	#	RV [km/s]	$v \sin i$ [km/s]	Multiplicity	Note
BD+18 2797	1467-0355		9.39	M5		1	-4.2			XZ Boo, LPV-known
HD 124569	1472-0535		7.48	M0III		1	-5.5			LPV-new
	1472-0811		9.71		M3-4(G)	1	-36.9			LPV-new
HD 142536	0943-0295		8.10	K5		3	26.4			LPV-new
HD 142639	0943-0347		6.89	K0	K0(G)	3	10.8			LPV-new
HD 149758	5636-0050		8.31	K5		2	-51.3			LPV-new
HD 149977	5636-0153		9.02	K5	K0(G)	2	-39.8			LPV-new
HD 175922	1051-1124		7.21	A _m	F5	1	-59.7			spots-new
HD 176285	1051-1430		9.51	F8	F4:	2	-14.6			
HD 176948	5136-0659		8.10	K2	K3(G)	2	6.7			LPV-known
HD 176949	5140-2635		8.88	G5	G2(G)	2	12.4			LPV-new
BD-05 4861	5136-1743		10.20	F5	F4-5	3	-26.5	112		V Algol, no-eclipses
HD 345479	1628-2407		11.30	F2	F3	1			SB2	gDor-new
HD 345478	1628-2345		10.51	F0	F4	1	26.2	119		dScut-new
HD 345467	2141-0566		8.22	M2		1	-2.1			LPV-new
HD 345468	2141-0922		9.55	F0	F4-5	5			SB2?	in vis. binary, gDor-new
HD 345444	2141-2482		10.53	A3	F2	1	-16.8	114		dScut-new
	2141-1146		9.69		K0:	2	-26.1			LPV-new
HD 345455	2141-0526		10.48	B9		2	-5.6			eclBin-new
HD 345459	2141-0916		7.58	K0	K1(G)	3	7.8			LPV-new
BD+08 4628	1105-0367		9.33	K2		1	5.2			LPV-new
BD+08 4631	1109-2546		9.35	G0	G8-9	1	-33.7			LPV-new
BD+18 4914	1688-1766		10.60	F5	F5-6	1	-38.5	40		gDor-dScut-known
BD+18 4923	1688-0773		8.72	K2		1	9.7			LPV-new
HD 209775	1684-1373		7.59	F0	F5-6	1	1.1	85		dScut-known
HD 217423	3216-2213		8.71	F0	F0	1	10.2	191		dScut-gDor-new

Explanation of columns: HD/BD/GSC – catalog identifications; Cluster membership; V – visual magnitude; Sp. (publ.) – published spectral type; Sp. (BF) – spectral type estimated from the BF strength; # – number of observed spectra; RV – the mean radial velocity; $v \sin i$ – projected rotational velocity; Multiplicity – as observed; Photometric variability type according to MOST observations. The abbreviations used: dScut – δ Sct pulsations, eclBin – eclipsing binary, gDor – γ Dor pulsations, LPV – Long-Period Variable, PMS – pre-main sequence variable, RGpuls – pulsating red giant. Few constant stars were also observed (given without variability type)

Notes on individual systems:

HD 965 – inconsistent rotational velocity: while our spectra indicate $v \sin i < 20 \text{ km s}^{-1}$, Uesugi & Fukuda (1970) give $v \sin i = 90 \text{ km s}^{-1}$

HD 9014 – SB2 composed of rapidly rotating F-type component with $v \sin i = 212 \pm 3 \text{ km s}^{-1}$ and slowly rotating much fainter component ($L_2/L_1 \approx 0.04$) with projected rotational rate below our spectral resolution

GSC 2816-2176 – according to Daniel et al. (1994) it is SB1 system. Our spectra, however, show two components of similar brightness

HD 32704 – relative intensity of components in BFs was found to significantly vary

HD 46180A – similar to HD 9014, rapidly rotating component has $v \sin i = 125 \pm 2 \text{ km s}^{-1}$, ratio of intensities of components is $L_2/L_1 = 0.11 \pm 0.03$

HD 61199 – our BF is blend of three components, separate RVs not determined

GSC 1472-0811 – TiO absorption strength indicates M3-4III spectral type

GCVS designations: HD 6242 = CQ Psc, HD 9289 = BW Cet, HD 258857 = V848 Mon, GSC 0814-0889 = ES Cnc, BD+18 2797 = XZ Boo, HD 209775 = V377 Peg

in a spectral window of about $240 \text{ \AA}$ around the Mg I triplet (5167, 5173 and $5184 \text{ \AA}$) with an effective resolving power of $R = 12,000 - 14,000$. The diffraction grating with 2160 lines/mm was used with the sampling of $0.117 \text{ \AA/pixel}$.

The region of the magnesium triplet has been used throughout the extensive short-period binary program (see Rucinski et al. (2008) for the most recently published paper of the DDO series) and we continued using it for this survey. The Mg I triplet feature serves exceedingly well for majority of late spectral types. Because of size constraints, short-period binaries with orbital periods shorter than one day which were observed during the binary program were practically all of late spectral types. For stars earlier than A0, however, the magnesium triplet becomes very weak and practically unusable for extraction of any radial velocity and broadening information. Even for mid-A type stars, metallic lines are weak and a high S/N is required. Therefore, several spectra of stars earlier than about A1 were found to be unusable and thus discarded from this survey. Three targets,

HD 31919, HD 32884 and HD 45972, which were included in the survey and assumed to have sufficiently late spectral types for the Mg I observations, were found later to be too early; they have been omitted from the tables.

One-dimensional spectra were extracted by the usual procedures within the IRAF environment¹ after the bias and flat-field photometric correction of the original frames. Cosmic-ray trails were removed using a program of Pych (2004).

The current survey was limited to objects north of -15° , and stars brighter than $V = 11.5$. The data were obtained between 2007 November 13 and 2008 June 19. Because of the observatory closure, most of the targets were observed only once or twice and the goals of the survey were

¹ IRAF is distributed by the National Optical Astronomy Observatories, which are operated by the Association of Universities for Research in Astronomy, Inc., under a cooperative agreement with the NSF.

Table 2. Parameters of the template stars used to extract broadening functions and to define the radial velocity system. HD 107966 was regularly observed, but due to its high rotational velocity it was not finally used as a template. The source of metallicity (always given relative to that of the Sun) is given in the last column.

No.	Name	RV [km/s]	$v \sin i$ [km/s]	sp.	[Fe/H]	Ref.
0	HD 72660	2.8	9	A1V	0.34	Lemke (1989)
1	HD 107966	1.3	55	A3V		
2	HD 128167	0.2	5	F2V	-0.41	Edvardsson et al. (1993)
3	HD 693	14.4	8	F5V	-0.38	Edvardsson et al. (1993)
4	HD 102870	4.6	3	F8.5IV-V	0.13	Edvardsson et al. (1993)
5	HD 114710	6.1		G0V	0.03	Edvardsson et al. (1993)
6	HD 23169	17.4		G2V		
7	HD 32923	20.3		G4V	-0.20	Hearnshaw (1972)
8	HD 65583	13.2		G8V	-0.70	Sopubiran et al. (2008)
9	HD 185144	26.7	8	K0V	-0.21	Sopubiran et al. (2008)
10	HD 3765	-62.3		K2V	0.10	Sopubiran et al. (2008)
11	HD 160964	-28.4		K4V		
12	HD 79210	11.1		M0V		

only partially fulfilled. In total, 241 spectra of 103 MOST targets (96 of them being photometrically variable) were obtained. An overview of the systems and their observations is given in Table 1. The dates of all spectroscopic observations are given in Table 3 listing the mean observed RVs for single and multiple stars, respectively. Selected BF of spectroscopic binaries and triple systems are shown in Fig. 1.

In addition to the program stars, several template spectra were observed (see Table 2) to be used in the broadening function determination (see the next Section 3). The templates were selected to be of stars with low projected rotational velocities (especially for A and F spectral types), known spectral types and constant radial velocities (hereafter RVs). Our template sequence is sufficiently continuous for the broadening function applications except for a gap between A1V (HD 72660) and F2V (HD 128167), where we could not find any suitable templates having a low projected RV. Several RV standards were also observed to ascertain their reliability within our RV system.

3 ANALYSIS OF SPECTRA

The observations were analyzed using the technique of broadening-functions (BF), described in Rucinski (1992, 2002), using spectra of slowly rotating stars of similar spectral type as templates. The directly extracted BFs were smoothed by a convolution with the Gaussian function to match the spectral resolution of the spectrograph. Smoothing with $\sigma = 1.5$ pixels (10 km s^{-1}) was found to be appropriate for most BFs; a few BFs which were very noisy were smoothed using $\sigma = 2$ pixels.

Because the observing time available for the observations was rather limited, mainly due to poor weather at DDO during the time of the survey and – later – due to the observatory closure, no low-resolution classification spectra were taken for direct spectral type and luminosity class estimates, as had been originally planned. The spectral types were, however, determined by finding a template (listed in Table 3) matching the observed spectrum best. For the best fitting template the integrated BF strength is expected to

be unity. This rough spectral-type estimate is possible because the wavelength range contains only metallic lines and their strength is increasing with decreasing temperature to about K5 spectral type where molecular bands start to be prominent. Therefore using progressively later templates for a given spectrum results in decreasing BF integral. For few stars we either could not achieve a good template match to the spectrum or the BF strength was not decreasing monotonically with progressively later-type templates. This typically occurred for stars later than K2–5 or B and A stars (some of which are probably chemically peculiar). For those stars, the spectral-type estimate based on the BF strength is not given (see Table 1). Except for the dwarf templates, we observed a few giants covering the range of spectral types of G8III to M3III. We noticed that using a dwarf template for the spectrum of a giant results in a significant depression around the resulting BF base. This could be used for a very approximate guess at the luminosity class. For targets showing such a depression which strongly indicates a luminosity class II–III, the spectral type in Table 1 was augmented by “(G)” (giant). In any case the spectral-type estimates should be regarded as very approximate because the BF strength is expected to depend not only on the temperature but also on the metallicity.

Determination of $v \sin i$ for low projected rotational velocities is limited by the spectral resolution ($R = 12\,000 - 15\,000$) setting a lower limit at about $15 - 20 \text{ km s}^{-1}$. The Gaussian profiles were fitted to BFs of slowly rotating stars with $v \sin i < 30 \text{ km s}^{-1}$ to determine RV; for those stars only the RV and estimated spectral type is given. For rapidly-rotating stars, the RVs and projected rotational velocities, $v \sin i$, were determined by fitting rotational profiles to the extracted broadening functions, as described in Pribulla et al. (2006). Because the shape of the theoretical rotational profile depends (weakly) on the limb darkening in the $5184 \text{ \AA}$ spectral window, the limb darkening coefficient was adopted from extensive tables of van Hamme (1993) according to the estimated spectral type.

Precision of the RV measurements is defined by random errors (influenced mainly by the S/N ratio of spectra,

Table 3. Radial velocities and their standard errors for all single stars, SB1 and SB2 systems and triple system ES Cnc (the full table is available only in electronic form, here RVs are given for first three SB2 systems and one single star). The column denoted as “Temp.” gives the template number used to extract BF and determine the RVs referring, as in the first column of Table 2. The components were designated according to observed relative brightness with the component 1 assumed being always the brighter one. The RV for SB2 systems is given if the components could be separated only. Type of the function to fit to BFs is given either as G (Gaussian) or R (rotational profile) in the columns denoted “Fit”. Radial velocity is not given for observations of SB2 or triple systems leading to inseparable peaks in BFs.

HJD 2 400 000+	Temp.	RV ₁ [km/s]	σ_1 [km/s]	Fit	RV ₂ [km/s]	σ_2 [km/s]	Fit
HD 9014							
54422.60716	2	−2.81	1.49	R	−3.11	1.05	G
54490.46861	2	−11.73	1.58	R	21.96	1.24	G
54494.46390	2	−14.78	1.71	R	23.41	1.11	G
54504.47860	2	−17.03	2.03	R	26.78	1.24	G
GSC 2816 2276							
54422.65196	3	36.86	0.58	G	−15.42	1.44	G
54481.66400	3	34.28	1.29	G	−14.58	2.76	G
54497.51571	3	−8.72	1.47	G	30.66	5.15	G
GSC 2816 2273							
54418.63188	3	−6.61	0.27	G	55.07	1.69	G
54490.49523	3	28.42	0.28	G	−20.29	0.73	G
HD 965							
54418.50574	2	0.77	0.39	G			
54418.53590	2	0.59	0.38	G			
54418.54755	2	0.82	0.39	G			




Figure 1. Selected broadening functions of SB2 and triple systems observed at DDO.

width and intensity of components in the extracted BFs) and by systematic effects of spectrograph flexure which are rather hard to quantify. Our observations did not utilize a iodine cell and telluric spectral lines were not present in our spectral window so no simple, external checks on the RV zero-point were available. We attempted to assess the stability of our RV system by using multiple observations of bright standard stars. For such stars having at least 5 spectra, the first spectrum of a given star was used as a template and the RVs were then determined fitting the (obviously very narrow) Gaussian function to the extracted BFs for the remaining spectra. As expected, the scatter within such samples depends on the spectral type of the standard and decreases for late-type standards with large numbers of strong lines in their spectra. Along the spectral sequence, we obtained: HD 97633 – A2V, $\sigma = 1.13 \text{ km s}^{-1}$, HD 128167 – F2V, $\sigma = 0.91 \text{ km s}^{-1}$, HD 102870 – F8.5IV-V, $\sigma = 0.84 \text{ km s}^{-1}$, and HD 144579 – G8V, $\sigma = 0.59 \text{ km s}^{-1}$. Hence, one can expect systematic deviations of individual RVs to be at the level of typically less than about $1.0 - 1.2 \text{ km s}^{-1}$. This effectively means that for measurements with formal standard errors given in Table 3 smaller than about $0.5 - 1.0 \text{ km s}^{-1}$, the systematic errors dominate in the error budget.

The resulting RVs for most observed systems are given in Table 3 (not given for binaries and multiples with complicated blends of components). With many targets observed just once, we cannot exclude a possibility that we missed several SB1 systems or SB2 systems observed close to their spectroscopic conjunctions.

4 BINARY AND MULTIPLE SYSTEMS

Spectroscopic observations showed that part of the targets are double or triple systems (see Fig. 1). RVs of individual components in these cases were determined by the same procedures as described in Pribulla et al. (2006). In the case of SB2 systems, double rotational or Gaussian profiles were fitted to the components, depending on the rotational broadening of the lines; the fitted functions are indicated in Table 3. For the triple system ES Cnc, the third (slowly-rotating) component was first removed after a multi-component Gaussian model was fit to the observed BFs; the profiles of the binary components were then approximated by rotational profiles. For the triple systems HD 61199 and HD 75638, where components are always blended, no individual RVs could be determined.

The present survey has led to detection of 7 new spectroscopic binaries: HD 9014 (GSC 5276-0313), HD 32704 (GSC 4758-0066), HD 46105 (GSC 0158-2674), HD 46180 (GSC 0154-2360) where both visual components are SB2, GSC 0814-0323, HD 345468 (GSC 2141-0922; components always blended), and HD 345479 (GSC 1628-2407; a marginal detection). Prior to the present spectroscopic survey, MOST photometry showed that HD 46180, and GSC 0814-0323 contain eclipsing binaries.

All spectroscopic binaries in NGC 752 observed within this survey were already detected by Daniel et al. (1994). This applies to GSC 2816-2068, GSC 2816-2176, GSC 2816-2273, and GSC 2816-2276. Unfortunately, the authors did not give any details regarding the orbital periods or preliminary spectroscopic orbits. The most interesting and impor-

tant is GSC 2816-2068, which was found to be an eclipsing binary with $P = 15.52$ days with eclipses about 0.11 mag deep. Four systems, GSC 0814-0323, HD 73709, HD 46180, and HD 75638 deserve special attention (see the following subsections). Notes for other systems are given below the main table (Table 1).

The spectroscopic elements (case of GSC 814-323 and HD 73709) were determined by the differential corrections optimization.²

4.1 GSC 0814-0323

The only newly discovered SB2 (and a eclipsing binary discovered by MOST) for which we have enough data to obtain a preliminary spectroscopic orbit is GSC 0814-0323, located in the sky close to the open cluster M67. The ratio of component luminosities, as estimated from the BFs when the components are split, is about $L_2/L_1 \sim 0.29$. With orbital period as long as 68 days, and with the obvious presence of eclipses, the orbital-plane inclination must be very close to 90° so that the projected masses of components, $M_1 = 1.39 \pm 0.08 M_\odot$ and $M_2 = 0.96 \pm 0.06 M_\odot$ must be close to the true ones. The predicted spectroscopic conjunction, HJD 2453547.35 ± 0.86 , occurs 8 days earlier than predicted by an eclipse observed by MOST and using the spectroscopic period. There is a single radial velocity measurement of GSC 0814-0323 at HJD 2441321.88 published by Mathieu et al. (1986), giving $RV = 12.6 \pm 0.5 \text{ km s}^{-1}$. The measurement probably refers to the brighter component of the visual pair and hence to the primary component of the eclipsing system. The light curve of GSC 0814-0323 was discussed in a paper dedicated to MOST photometry of M67 field (Pribulla et al. 2008).

4.2 HD 73709

HD 73709 was found to be SB1 by Abt & Willmarth (1999) who determined the following orbital elements: $P = 7.2204(1)$ days, $V_0 = 34.6(1) \text{ km s}^{-1}$ and $K = 31.2(2) \text{ km s}^{-1}$, based on 30 RV measurements over 3 years. MOST photometry showed that the star is an eclipsing binary with $P = 7.22$ days, which is compatible with the spectroscopic determination. The system is a high-probability member of the Praesepe cluster. Our new RVs were combined with those of Abt & Willmarth (1999) to refine the orbital elements. Since the orbital eccentricity was found to be insignificant, we assumed the circular orbit. All RVs and the combined solution are shown in Fig. 2.

4.3 HD 46180

HD 46180 (GSC 0154-2360) is close visual pair in NGC 2244 composed of components of $V = 9.1 + 9.3$ at $\theta = 82^\circ$, and $\rho = 5.2''$ (Mason et al. 2001). The stars were oriented almost along the spectrograph slit which enabled us to simultaneously obtain spectra of either of the components on nights with the best seeing. Our spectroscopy shows that each of the visual components is a binary star so that HD 46180

² Unpublished code *ELEMDR* of the first author




Figure 2. The radial velocity orbits for the SB2 system GSC 0814-0323 (top) and the SB1 system HD 73709 (bottom). One radial-velocity determination for the former star (plotted by an open symbols) was not used in the computation of the orbit. The new RVs of HD 73709 were combined with those published in Abt & Willmarth (1999) to refine the orbital period (see the text). The resulting preliminary orbital elements are listed in the respective panels.

is a quadruple system. Unfortunately, we were able to obtain only 10 spectra of the brighter visual component and 4 of the fainter one (on other nights it was blended with the brighter component), which has prevented us to determine the respective orbital periods. MOST satellite photometry refers to the whole quadruple system and shows that one of the binaries is an eclipsing one with the period of $P = 3.09$ days and the photometric amplitude of only 0.02 mag. The present data indicate that the brighter component of the visual pair is the variable one.

4.4 HD 75638

HD 75638 (GSC 0814-0601) is a visual triple system WDS 08515+1208 consisting of components A ($V = 8.28$), B ($V = 10.17$) and C ($V = 11.8$). Our spectroscopy refers to the pair AB, $\rho = 1.4$ arcsec, which could not be separated on the DDO spectrograph slit. The pair AB was suspected to be an SB3 system in the survey of Nordstroem et al. (1997). Their analysis of 13 spectra showed two measurable components with $v_1 \sin i = 80 \text{ km s}^{-1}$ and $v_2 \sin i = 10 \text{ km s}^{-1}$, temperatures $T_1 = 7000K$, and $T_2 = 6750K$, and a luminosity ratio of 0.10. A preliminary orbit for the fainter component

was given: $P = 5.8167(9)$ days, $V_0 = 11.05(91) \text{ km s}^{-1}$, $K_1 = 28.9(12) \text{ km s}^{-1}$, $e = 0.083(56)$, and $\omega = 322(28)^\circ$. The spectroscopic orbit most likely corresponds to the motion of the blend of a faint pair seen as component B.

The MOST photometry showed very shallow ($\Delta m = 0.013$) eclipses giving a preliminary mid-eclipse ephemeris, $\text{HJD } 2454145.85 + 5.81 \times E$, in good accord the spectroscopic result of Nordstroem et al. (1997) (see Pribulla et al. (2008)). Our spectroscopic results are related to the AB components. The broadening functions (Fig. 1) show the brighter, rapidly-rotating component ($v \sin i \approx 85 \text{ km s}^{-1}$) to be stationary in the radial velocity; this star is most likely the A component of the visual pair. The fainter component B occasionally splits into two. All three components are always blended, suggesting that a high-resolution spectroscopy will be necessary to reliably define orbits of both components.

5 CONCLUSIONS

The medium dispersion spectroscopic survey of the variable stars discovered by the MOST satellite was planned to include some 150 targets observed several times to fully ascertain types of variability. This has not been possible because of the DDO observatory closure. We publish here radial velocities and spectral type estimates for 103 MOST targets, 96 being variables with 83 of them being new MOST detections. For 31 targets, we give the projected rotational velocities $v \sin i$.

The survey resulted in the discovery of seven new spectroscopic binaries. For two of these binaries (HD 73709, and GSC 0814-0323), which were found to show eclipses by the MOST satellite photometry, preliminary spectroscopic orbits were determined. HD 46180 was found to be composed of two binary stars forming, very probably, a physical quadruple system. According to the MOST photometry, one of the binaries is eclipsing. The newly discovered spectroscopic binaries will require further spectroscopy to refine their orbital periods and the remaining orbital parameters.

ACKNOWLEDGMENTS

This study has been funded by the Canadian Space Agency Space Enhancement Program (SSEP) with TP holding a Post-Doctoral Fellowship position at the University of Toronto. The Natural Sciences and Engineering Research Council of Canada (NSERC) supports the research of SMR. Support from the Polish Science Committee (KBN grants PO3D 006 22 and PO3D 003 24) to WO is acknowledged with gratitude.

The authors would like to thank the telescope operators at DDO, Heide de Bond, Toomas Karmo and Archie de Ridder, for their help.

The research made use of the SIMBAD database, operated at the CDS, Strasbourg, France and accessible through the Canadian Astronomy Data Centre, which is operated by the Herzberg Institute of Astrophysics, National Research Council of Canada. This research made also use of the Washington Double Star (WDS) Catalog maintained at the U.S. Naval Observatory.

REFERENCES

- Abt, H.A., Willmarth, D.W. 1999, ApJ, 521, 682
- Daniel, S.A., Latham, D.W., Mathieu, R.D., Twarog, B.A. 1994, PASP, 106, 281
- Edvardsson, B., Andersen, J., Gustafsson, B., Lambert, D.L., Nissen, P.E., Tomkin, J. 1993, A&A, 275, 101
- Hearnshaw, J.B. 1972, Mem. R. Astron. Soc., 77, 55
- Hearnshaw, J.B. 1976, A&A, 51, 71
- Lemke, M. 1989, A&A, 225, 125
- Mason, B.D., Wycoff, G.L., Hartkopf, W.I., Douglass, G.G., & Worley, C.E. 2001, AJ, 122, 3466 (WDS)
- Mathieu R.D., Latham D.W., Griffin R.F., Gunn J.E. 1986, AJ, 92, 1100
- Matthews J.M., Kusching R., Guenther D.B., Walker G.A.H., Moffat A.F.J., Rucinski S.M., Sasselov D., Weiss W.W. 2004, Nature, 430, 51
- Nordstrom B., Stefanik R.P., Latham D.W., Andersen J., 1997, A&AS, 126, 21
- Pribulla, T., Rucinski, S.M., Lu, W., Mochnacki, S.W., Conidis, G., DeBond, H., Thomson, J.R., Pych, W. Blake, R.M., Ogloza, W., Siwak, M. 2006, AJ, 132, 769
- Pribulla, T., Rucinski, S.M., Matthews, J.M., Kuschnig, R., Rowe, J.F., Guenther, D.B., Moffat, A.F.J., Sasselov, D., Walker, G.A.H., Weiss, W.W. 2008, MNRAS, in press
- Pych W. 2004, PASP, 116, 148
- Rucinski S.M. 1992, AJ, 104, 1968
- Rucinski S.M. 2002, AJ, 124, 1746
- Rucinski, S.M., Pribulla, T., Mochnacki, S., Liokumovich, E., Lu, W., DeBond, H., de Ridder, A., Karmo, T., Rock, M., Thomson, J.R., Ogloza, W., Kaminski, K., Ligeza, P. 2008, AJ, 136, 586
- Soubiran, C., Bienayme, O., Mishenina, T.V., Kovtyukh, V.V. 2008, A&A, 480, 91
- Uesugi, A., Fukuda, I. 1970, Contr. Astroph. Kwasan Obs. Univ. Kyoto No. 189
- Walker G., Matthews J., Kuschnig R., Johnson R., Rucinski S., Pazder J., Burley G., Walker A., Skaret K., Zee R., Grocott S., Carroll K., Sinclair P., Sturgeon D., Harron J. 2003, PASP, 115, 1023
- van Hamme W. 1993, AJ, 106, 2096